

# ON THE EXPONENTIAL BONDAGE NUMBER OF A GRAPH

VECDİ AYTAÇ, TUFAN TURACI

**ABSTRACT.** The domination number is an important vulnerability parameter that has become one of the most widely studied topics in graph theory. The bondage number is related to the domination number, and also the most often studied property of vulnerability of communication networks. Recently, Dankelmann et al. have defined the exponential domination number in [17]. We investigate a refinement that involves the exponential bondage number of this parameter. Let  $G = (V(G), E(G))$  be a simple graph. The exponential bondage number, denoted by  $b_{exp}(G)$ , is defined by  $b_{exp}(G) = \min\{|B_e| : B_e \subseteq E(G), \gamma_e(G - B_e) > \gamma_e(G)\}$ , where  $\gamma_e(G)$  is the exponential domination number of  $G$ . In this paper, the above mentioned new parameter is defined and examined. Then upper bounds, lower bounds and exact formulas are obtained for any graph  $G$ . Finally, the exact values have been determined for some well-known graph families.

**Mathematics Subject Classification (2010):** 05C40, 05C69, 68M10, 68R10.

**Key words:** Graph vulnerability; Connectivity; Domination number; Bondage number; Exponential domination number; Exponential bondage number.

*Article history:*

Received 5 January 2018

Accepted 21 May 2018

## 1. INTRODUCTION

Graph theory has seen an explosive growth due to interaction with areas like computer science, operation research, etc. Especially, it has become one of the most powerful mathematical tools in the analysis and study of the architecture of a network. A network is described as an undirected and unweighted graph in which vertices represent the processing and edges represent the communication channel between them [8, 9, 13].

It is known that communication systems are often exposed to failures and attacks. So vulnerability of the network topology is a key aspect in the design of computer networks. The vulnerability value of a communication network shows the resistance of the network after the disruption of some centers or connection lines until a communication breakdown. In the literature, various measures have been defined to measure the robustness of network and a variety of graph theoretic parameters have been used to derive formulas to calculate network vulnerability. Graph vulnerability relates to the study of graph when some of its elements (vertices or edges) are removed. The measures of graph vulnerability are usually invariants that measure how a deletion of one or more network elements changes properties of the network. The best known measure of reliability of a graph is its connectivity. The vertex (edge) connectivity is defined to be the minimum number of vertices (edges) whose deletion results in a disconnected or trivial graph [8]. Then toughness [25], integrity [5], domination number [6, 19], bondage number [3, 4, 7, 10], reinforcement number [12] etc. have been proposed for measuring the vulnerability of networks. Recently, some average vulnerability parameters such as average lower independence number [2, 14, 24], average lower domination

number [14, 23], average connectivity number [15], average lower bondage number [20] and average lower reinforcement number [21] have been defined.

Let  $G = (V(G), E(G))$  be a simple undirected graph of order  $n$ . We begin by recalling some standard definitions that we need throughout this paper. For any vertex  $v \in V(G)$ , the *open neighborhood* of  $v$  is  $N_G(v) = \{u \in V(G) | uv \in E(G)\}$  and *closed neighborhood* of  $v$  is  $N_G[v] = N_G(v) \cup \{v\}$ . The degree of  $v$  in  $G$  denoted by  $deg(v)$ , is the size of its open neighborhood. A vertex  $v$  is said to be pendant if  $deg(v) = 1$  [6, 19]. A vertex  $u$  is called support vertex if  $u$  is adjacent to a pendant vertex. The graph  $G$  is called  $r$ -regular graph if  $deg(v) = r$  for every vertex  $v \in V(G)$ . The complement  $\overline{G}$  of a graph  $G$  has  $V(G)$  as its vertex sets, but two vertex are adjacent in  $\overline{G}$  if only if they are not adjacent in  $G$ . The *distance*  $d(u, v)$  between two vertices  $u$  and  $v$  in  $G$  is the length of a shortest path between them. The *diameter* of  $G$ , denoted by  $diam(G)$  is the largest distance between two vertices in  $V(G)$  [6, 19]. A cycle passing through all the vertices of a graph is called a *Hamiltonian cycle*. A graph containing a Hamiltonian cycle is called a *Hamiltonian graph*, and also if there exist a path  $P$  in the graph  $G$  such that  $V(P) = V(G)$ , then  $G$  is called *semihamiltonian* [19]. A *vertex-transitive graph* is a graph such that every pair of vertices is equivalent under some element of its automorphism group [26]. A set of pairwise independent edges of  $G$  is called a *matching* in  $G$ , while a matching of maximum cardinality is a maximum matching in  $G$  [19]. A *vertex cover* of a graph  $G$  can also more simply be thought of as a set  $S$  of vertices of  $G$  such that every edge of  $G$  has at least one of member of  $S$  as an endpoint. The vertex set of a graph is therefore always a vertex cover. The smallest possible vertex cover for a given graph  $G$  is known as a minimum vertex cover, and its size is called the *vertex cover number*, denoted by  $\tau(G)$ . An *edge cover* is a subset of edges defined similarly to the vertex cover, namely a collection of graph edges such that the union of edge endpoints corresponds to the entire vertex set of the graph. Therefore, only graphs with no isolated points have an edge cover. An edge cover having the smallest possible number of edges for a given graph is known as a *edge cover number*, denoted by  $\tau'(G)$  [18]. The smallest integer not less than  $x$  is denoted by  $\lceil x \rceil$ . A set  $S \subseteq V(G)$  is a *dominating set* if every vertex in  $V(G) - S$  is adjacent to at least one vertex in  $S$ . The minimum cardinality taken over all dominating sets of  $G$  is called the *domination number* of  $G$  is denoted by  $\gamma(G)$  [6, 19].

There are different application of domination problems. For instance, dominating sets in graphs are natural models for facility location problems in operations research [19] or domination number is the one of the most important vulnerability parameter for networks [19, 23]. When investigating the domination number of a given graph  $G$ , one may want to know the answer of the following question: How does the domination number increases in a graph  $G$ ? or How many edges need to be added to decrease the domination number of the original graph? One of the vulnerability parameters known as *bondage number* in a graph  $G$  answers the former question. The bondage number  $b(G)$  was introduced by Fink et al. [10] and is defined as follows:

$$b(G) = \min\{|B| : B \subseteq E(G), \gamma(G - B) > \gamma(G)\}.$$

We call such an edge set  $B$  that  $\gamma(G - B) > \gamma(G)$  the *bondage set* and the minimum one the *minimum bondage set*. If  $E(G) = \emptyset$ , then we say that  $b(G) = \infty$ .

The *reinforcement number* in a graph  $G$  answers the latter question. The reinforcement number  $r(G)$  was introduced by Kok et al.[12] and is defined as follows:

$$r(G) = \min\{|R| : R \subseteq E(\overline{G}), \gamma(G) > \gamma(G + R)\}.$$

We call such an edge set  $R \subseteq E(\overline{G})$  a reinforcement set, if  $\gamma(G) > \gamma(G + R)$ .

In 2009, Dankelmann introduced the concept of *exponential domination*[17]. This new parameter is closely in relation with distance of each pair of vertices. The exponential domination number is the theoretical vulnerability parameters for a network that is represented by a graph. An exponential dominating set of graph  $G$  is subset  $S \subseteq V(G)$  such that  $\sum_{v \in S} (1/2)^{\overline{d}(u,v)-1} \geq 1, \forall v \in V(G)$ , where  $\overline{d}(u, v)$  is the length of a shortest path in  $\langle V(G) - (S - \{u\}) \rangle$  if such a path exist, and  $\infty$  otherwise. The

minimum exponential domination number,  $\gamma_e(G)$  is the smallest cardinality of an exponential dominating set. We call such an edge set is a minimum exponential set or a  $\gamma_e$ -set.

Our aim in this paper is to define a new vulnerability parameter, so called exponential bondage number. In Section 2, some well-known basic results are given for exponential domination number and bondage number. In Section 3, we define a new parameter namely the exponential bondage number denoted by  $b_{exp}(G)$ . In Section 4, we determine upper bounds, lower bounds and exact solutions of the exponential bondage number for any graph  $G$ . Finally, the exponential bondage numbers of the popular well-known graphs are computed in Section 5.

## 2. BASIC RESULTS

In this section some well-known basic results are given with regard to exponential domination number and bondage number.

**Theorem 2.1.** [17] *The exponential domination number of*

- (a) *the path graph  $P_n$  of order  $n \geq 2$  is  $\gamma_e(P_n) = \lceil \frac{n+1}{4} \rceil$ .*
- (b) *the cycle graph  $C_n$  of order  $n \geq 4$  is  $\gamma_e(C_n) = \begin{cases} 2 & , \text{if } n = 4; \\ \lceil \frac{n}{4} \rceil & , \text{if } n \neq 4. \end{cases}$*

**Theorem 2.2.** [17] *For every graph  $G$ ,  $\gamma_e(G) \leq \gamma(G)$ , and also  $\gamma_e(G) = 1$  if and only if  $\gamma(G) = 1$ .*

**Theorem 2.3.** [1] *Let  $G$  be any connected graph with  $n$  vertices and  $\exists v \in V(G)$  such that  $\deg(v) = n - 1$ . Then,  $\gamma_e(G) = 1$*

**Theorem 2.4.** [10] *If  $G$  is a connected graph of order  $n \geq 2$ , then  $b(G) \leq n - \gamma(G) + 1$ .*

**Theorem 2.5.** [10] *The bondage number of*

- (a) *the path graph  $P_n$  of order  $n \geq 2$  is  $b(P_n) = \begin{cases} 2 & , \text{if } n \equiv 1 \pmod{3}; \\ 1 & , \text{otherwise.} \end{cases}$*
- (b) *the cycle graph  $C_n$  of order  $n \geq 3$  is  $b(C_n) = \begin{cases} 3 & , \text{if } n \equiv 1 \pmod{3}; \\ 2 & , \text{otherwise.} \end{cases}$*
- (c) *the complete graph  $K_n$  of order  $n \geq 2$  is  $b(K_n) = \lceil \frac{n}{2} \rceil$ .*
- (d) *the star graph  $S_n$  of order  $n \geq 3$  is  $b(S_n) = 1$ .*

**Theorem 2.6.** [22] *If  $G$  is a nonempty graph with a unique minimum dominating set, then  $b(G) = 1$ .*

**Theorem 2.7.** [11] *Let  $G$  be a vertex-transitive graph. Then,  $b(G) \geq \lceil \frac{n}{2\gamma(G)} \rceil$ .*

## 3. THE EXPONENTIAL BONDAGE NUMBER

In this section, we introduce a new graph theoretical parameter: the *exponential bondage number* and it is defined as:

$$b_{exp}(G) = \min\{|B_e| : B_e \subseteq E(G), \gamma_e(G - B_e) > \gamma_e(G)\},$$

where  $\gamma_e(G)$  is the exponential domination number of the graph  $G$ . We call such an edge set  $B_e$  that  $\gamma_e(G - B_e) > \gamma_e(G)$  the *exponential bondage set* and the minimum one the *minimum exponential bondage set*.

If we think of a graph as a modeling of network, the exponential bondage number may be more sensitive than other measures of vulnerability such as connectivity, domination number, exponential domination number and bondage number for distinguish two graphs whose number the vertices and edges are the same. For example, consider two graphs  $G_1$  and  $G_2$  in Figure1, where  $|V(G_1)| = |V(G_2)| = 7$  and  $|E(G_1)| = |E(G_2)| = 8$ . They have not only equal connectivity but also equal domination number, exponential domination number and bondage number such as  $k(G_1) = k(G_2) = 1$ ,  $\gamma(G_1) = \gamma(G_2) = 2$ ,  $\gamma_e(G_1) = \gamma_e(G_2) = 2$  and  $b(G_1) = b(G_2) = 1$ . These values could be easily checked by readers. So, how can we distinguish between the graphs  $G_1$  and  $G_2$ ?




FIGURE 1. The graphs  $G_1$  and  $G_2$

When the exponential bondage numbers of these two graphs  $G_1$  and  $G_2$  are computed,  $b_{exp}(G_1) = 1$  and  $b_{exp}(G_2) = 2$  are obtained. The results could be checked by readers. Thus, the exponential bondage number may be used for distinguish between these two graphs  $G_1$  and  $G_2$ .

#### 4. UPPER BOUNDS, LOWER BOUNDS AND EXACT FORMULAS

**Theorem 4.1.** *If  $\gamma(G) = \gamma_e(G)$ , then  $b_{exp}(G) \geq b(G)$ .*

*Proof.* Let  $B_e$  be a minimum exponential bondage set of graph  $G$ . Then  $\gamma(G) = \gamma_e(G) \leq \gamma_e(G - B_e) \leq \gamma(G - B_e)$  by Theorem 2.2. Thus, we have  $b_{exp}(G) \geq b(G)$ .

The proof is completed.  $\square$

**Theorem 4.2.** *If  $G$  is a connected graph, then  $b_{exp}(G) \leq \delta(G)$ .*

*Proof.* To exponentially dominate all the vertices as the minimum number, first of all the vertex  $u$  with maximum degree should be taken to  $\gamma_e$ -set. That is, the vertex  $u$  belongs to a minimum exponential dominating set. Let  $D$  be  $\gamma_e$ -set for  $G$ . We know that  $u \in D$ . Suppose that the graph  $G$  has a vertex  $v$  with minimum degree,  $deg(v) = \delta(G)$ , that does not belong to  $D$ . Let  $E_v$  denote the set of edges incident with  $v$ . The minimum exponential dominating set for  $G - E_v$  is  $D + \{v\}$ . Hence  $\gamma_e(G - E_v) > \gamma_e(G)$  is obtained. As a result, we have  $b_{exp}(G) \leq |E_v| = deg(v) \leq \delta(G)$ .

The proof is completed.  $\square$

**Theorem 4.3.** *If  $G$  is a vertex-transitive graph and  $\gamma(G) = \gamma_e(G)$ , then*

$$b_{exp}(G) \geq 2 \lceil \frac{n}{2\gamma_e(G)} \rceil + \gamma_e(G) - (n + 1).$$

*Proof.* By Theorems 2.4 and 4.1, we have  $b_{exp}(G) \geq 2b(G) + \gamma_e(G) - (n + 1)$ . By using Theorem 2.7, it is trivial that

$$b_{exp}(G) \geq 2 \lceil \frac{n}{2\gamma_e(G)} \rceil + \gamma_e(G) - (n + 1).$$

The proof is completed.  $\square$

**Theorem 4.4.** *Let  $G$  be a connected graph of order  $n$ . If  $G$  includes only one pendant vertex, then  $b_{exp}(G) = 1$ .*

*Proof.* Let  $v$  be the pendant vertex of  $G$ . The removal of an edge  $e$  which is incident to  $v$  from the graph  $G$  leaves a graph  $G' \cong G - \{e\}$  consisting of two components. One of these is an isolated vertex and the other is connected graph  $H$  with  $n - 1$  vertices. That is,  $G' = H \cup \{v\}$ . Let  $D$  be  $\gamma_e$ -set for  $G$ . Since the set  $D$  must include the support vertex  $u$  which is adjacent to the vertex  $v$ , it is easy to see that  $\gamma_e(H) = \gamma_e(G)$ . Hence the  $\gamma_e$ -set for  $G'$  is  $D \cup \{v\}$ . Consequently, since  $\gamma_e(G') > \gamma_e(G)$  is obtained, the minimum exponential bondage set of  $G$  has an edge which is incident to vertex  $v$ . Thus, we have  $b_{exp}(G) = 1$ .

The proof is completed.  $\square$

## 5. THE EXPONENTIAL BONDAGE NUMBER OF SOME WELL-KNOWN GRAPHS

In this section we calculate the exponential bondage number of some well known graphs such as the path graph  $P_n$ , the cycle graph  $C_n$ , the complete graph  $K_n$ , the star graph  $S_n$  and the wheel graph  $W_n$ . Furthermore, we give some corollaries for any graph  $G$ .

**Theorem 5.1.** *The exponential bondage number of the path graph  $P_n$  of order ( $n \geq 2$ ) is given by  $b_{exp}(P_n) = 1$ .*

*Proof.* While we are calculating the exponential bondage number of the path graph  $P_n$ , we have four cases according to the number of vertices of  $P_n$ .

*Case 1.*  $n \equiv 0(\text{mod } 4)$ .

We know that  $\gamma_e(P_n) = \lceil \frac{n+1}{4} \rceil$  by Theorem 2.1. The removal of an edge from  $P_n$  leaves a graph  $H$  consisting of two paths  $P_{n_1}$  and  $P_{n_2}$ , where  $n_1 + n_2 = n$ . Then, one of the following statements is satisfied.

- $n_1 \equiv 1(\text{mod } 4)$  and  $n_2 \equiv 3(\text{mod } 4)$
- $n_1 \equiv n_2 \equiv 2(\text{mod } 4)$
- $n_1 \equiv n_2 \equiv 0(\text{mod } 4)$

In the first case,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \lceil \frac{n_1+1}{4} \rceil + \lceil \frac{n_2+1}{4} \rceil \\ &= \frac{n_1+3}{4} + \frac{n_2+1}{4} = \frac{n+4}{4} = \lceil \frac{n+1}{4} \rceil = \gamma_e(P_n). \end{aligned}$$

In the second case,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \lceil \frac{n_1+1}{4} \rceil + \lceil \frac{n_2+1}{4} \rceil \\ &= \frac{n_1+2}{4} + \frac{n_2+2}{4} = \frac{n+4}{4} = \lceil \frac{n+1}{4} \rceil = \gamma_e(P_n). \end{aligned}$$

In the last case,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \lceil \frac{n_1+1}{4} \rceil + \lceil \frac{n_2+1}{4} \rceil \\ &= \frac{n_1+4}{4} + \frac{n_2+4}{4} = \frac{n+8}{4} = \frac{n+4}{4} + 1 \\ &= \lceil \frac{n+1}{4} \rceil + 1 = \gamma_e(P_n) + 1 > \gamma_e(P_n). \end{aligned}$$

So, if the graph  $H$  is obtained by removing exactly one edge from  $P_n$  such that  $n_1 \equiv n_2 \equiv 0(\text{mod } 4)$ , then we get  $b_{exp}(P_n) = 1$ .

*Case 2.*  $n \equiv 1(\text{mod } 4)$ .

Let  $H$  be a graph obtained by the deletion of an edge of  $P_n$ . Then, we get either  $n_1 \equiv 1(\text{mod } 4)$  and  $n_2 \equiv 0(\text{mod } 4)$ , or  $n_1 \equiv 2(\text{mod } 4)$  and  $n_2 \equiv 3(\text{mod } 4)$ .

In the former case,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \lceil \frac{n_1+1}{4} \rceil + \lceil \frac{n_2+1}{4} \rceil \\ &= \frac{n_1+3}{4} + \frac{n_2+4}{4} = \frac{n+7}{4} = \frac{n+3}{4} + 1 \\ &= \lceil \frac{n+1}{4} \rceil + 1 = \gamma_e(P_n) + 1 > \gamma_e(P_n). \end{aligned}$$

In the latter case,

$$\gamma_e(H) = \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \lceil \frac{n_1+1}{4} \rceil + \lceil \frac{n_2+1}{4} \rceil$$

$$= \frac{n_1+2}{4} + \frac{n_2+1}{4} = \frac{n+3}{4} = \lceil \frac{n+1}{4} \rceil = \gamma_e(P_n).$$

If the graph  $H$  is obtained by removing one edge from  $P_n$  consists of an isolated vertex and a path of order  $n-1$ , then we get  $\gamma_e(H) > \gamma_e(P_n)$ . Consequently, we have  $b_{exp}(P_n) = 1$ .

*Case 3.*  $n \equiv 2(\text{mod } 4)$ .

Let  $H$  be a graph obtained by removing an edge from  $P_n$ . Then, one of the following statements is satisfied.

- $n_1 \equiv n_2 \equiv 1(\text{mod } 4)$
- $n_1 \equiv n_2 \equiv 3(\text{mod } 4)$
- $n_1 \equiv 2(\text{mod } 4)$  and  $n_2 \equiv 0(\text{mod } 4)$

In the first case,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \lceil \frac{n_1+1}{4} \rceil + \lceil \frac{n_2+1}{4} \rceil \\ &= \frac{n_1+3}{4} + \frac{n_2+3}{4} = \frac{n+6}{4} = \lceil \frac{n+2}{4} \rceil + 1 = \gamma_e(P_n) + 1 > \gamma_e(P_n). \end{aligned}$$

In the second case, it is easy to see that  $\gamma_e(H) = \gamma_e(P_n)$ .

In the last case,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \lceil \frac{n_1+1}{4} \rceil + \lceil \frac{n_2+1}{4} \rceil \\ &= \frac{n_1+2}{4} + \frac{n_2+4}{4} = \frac{n+6}{4} = \frac{n+2}{4} + 1 \\ &= \lceil \frac{n+1}{4} \rceil + 1 = \gamma_e(P_n) + 1 > \gamma_e(P_n). \end{aligned}$$

If an edge is removed from  $P_n$  such that either  $n_1 \equiv n_2 \equiv 1(\text{mod } 4)$  or,  $n_1 \equiv 2(\text{mod } 4)$  and  $n_2 \equiv 0(\text{mod } 4)$ , then  $\gamma_e(H) > \gamma_e(P_n)$ . Hence, we get  $b_{exp}(P_n) = 1$ .

*Case 4.*  $n \equiv 3(\text{mod } 4)$ .

Removing an edge from  $P_n$ , we have either  $n_1 \equiv 1(\text{mod } 4)$  and  $n_2 \equiv 2(\text{mod } 4)$ , or  $n_1 \equiv 0(\text{mod } 4)$  and  $n_2 \equiv 3(\text{mod } 4)$ .

In the former case,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \lceil \frac{n_1+1}{4} \rceil + \lceil \frac{n_2+1}{4} \rceil \\ &= \frac{n_1+3}{4} + \frac{n_2+2}{4} = \frac{n+5}{4} = \frac{n+1}{4} + 1 \\ &= \lceil \frac{n+1}{4} \rceil + 1 = \gamma_e(P_n) + 1 > \gamma_e(P_n). \end{aligned}$$

In the latter case,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \lceil \frac{n_1+1}{4} \rceil + \lceil \frac{n_2+1}{4} \rceil \\ &= \frac{n_1+4}{4} + \frac{n_2+1}{4} = \frac{n+5}{4} = \lceil \frac{n+1}{4} \rceil + 1 \\ &= \lceil \frac{n+1}{4} \rceil + 1 = \gamma_e(P_n) + 1 > \gamma_e(P_n). \end{aligned}$$

By the above both cases, we have  $\gamma_e(H) > \gamma_e(P_n)$ . Thus,  $b_{exp}(P_n) = 1$  is obtained.

By Cases 1, 2, 3 and 4, the exponential bondage number of the path graph of order  $n$  is  $b_{exp}(P_n) = 1$ .

The proof is completed.  $\square$

**Theorem 5.2.** *The exponential bondage number of the cycle graph  $C_n$  of order  $(n \geq 4)$  is given by*

$$b_{exp}(C_n) = \begin{cases} 1 & , \text{if } n \equiv 0(\text{mod } 4); \\ 2 & , \text{otherwise.} \end{cases}$$

*Proof.* We know that  $\gamma_e(C_n) = \lceil \frac{n}{4} \rceil$  for  $n \neq 4$  by Theorem 2.1. We have four cases depending on  $n$ .

*Case 1.*  $n \equiv 0(\text{mod } 4)$ .

The removal of one edge from the graph  $C_n$  leaves a graph  $H$  consisting of a path of order  $n$ . Thus,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_n) \\ &= \frac{n+4}{4} = \frac{n}{4} + 1 = \gamma_e(C_n) + 1 > \gamma_e(C_n). \end{aligned}$$

Since  $\gamma_e(H) > \gamma_e(C_n)$  is obtained, we get  $b_{exp}(C_n) = 1$ .

*Case 2.*  $n \equiv 1(\text{mod } 4)$ .

If the graph  $H$  is obtained by removing an edge from  $C_n$ , then the remaining graph is the path graph with  $n$  vertices. Since  $\gamma_e(H) = \gamma_e(P_n) = \frac{n+3}{4} = \gamma_e(C_n)$  for  $n > 4$ , we have  $b_{exp}(C_n) \geq 2$ .

If  $n \equiv 1(\text{mod } 4)$ , the removal of two edges from  $C_n$ , leaves a graph  $H$  consisting of two paths  $P_{n_1}$  and  $P_{n_2}$ , where  $n_1 + n_2 = n$ .

By using the Case 2 of the proof of Theorem 5.1, we obtain  $n_1 \equiv 1(\text{mod } 4)$  and  $n_2 \equiv 0(\text{mod } 4)$ . Thus,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \frac{n_1+3}{4} + \frac{n_2+4}{4} \\ &= \frac{n+7}{4} = \frac{n+3}{4} + 1 = \gamma_e(C_n) + 1 > \gamma_e(C_n). \end{aligned}$$

Since  $\gamma_e(H) > \gamma_e(C_n)$ , we have  $b_{exp}(C_n) = 2$ .

*Case 3.*  $n \equiv 2(\text{mod } 4)$ .

This case is very similar to the Case 2. By using the Case 3 of the proof of Theorem 5.1, we obtain directly either  $n_1 \equiv n_2 \equiv 1(\text{mod } 4)$  or,  $n_1 \equiv 2(\text{mod } 4)$  and  $n_2 \equiv 0(\text{mod } 4)$ .

In the former case,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \frac{n_1+3}{4} + \frac{n_2+3}{4} \\ &= \frac{n+6}{4} = \frac{n+2}{4} + 1 = \gamma_e(C_n) + 1 > \gamma_e(C_n). \end{aligned}$$

In the latter case,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \frac{n_1+2}{4} + \frac{n_2+4}{4} \\ &= \frac{n+6}{4} = \frac{n+2}{4} + 1 = \gamma_e(C_n) + 1 > \gamma_e(C_n). \end{aligned}$$

Since  $\gamma_e(H) > \gamma_e(C_n)$ , we have  $b_{exp}(C_n) = 2$ .

*Case 4.*  $n \equiv 3(\text{mod } 4)$ .

This case is also very similar to the Case 2. By using the Case 4 of the proof of Theorem 5.1, it is easy to see that we have either  $n_1 \equiv 1(\text{mod } 4)$  and  $n_2 \equiv 2(\text{mod } 4)$  or,  $n_1 \equiv 0(\text{mod } 4)$  and  $n_2 \equiv 3(\text{mod } 4)$ .

In the former case,

$$\begin{aligned} \gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \frac{n_1+3}{4} + \frac{n_2+2}{4} \\ &= \frac{n+5}{4} = \frac{n+1}{4} + 1 = \gamma_e(C_n) + 1 > \gamma_e(C_n). \end{aligned}$$

In the latter case,

$$\begin{aligned}\gamma_e(H) &= \gamma_e(P_{n_1}) + \gamma_e(P_{n_2}) = \frac{n_1+4}{4} + \frac{n_2+1}{4} \\ &= \frac{n+5}{4} = \frac{n+1}{4} + 1 = \gamma_e(C_n) + 1 > \gamma_e(C_n).\end{aligned}$$

Since  $\gamma_e(H) > \gamma_e(C_n)$ , we have  $b_{exp}(C_n) = 2$ .

Consequently, By Cases 1, 2, 3 and 4, the proof is completed.  $\square$

From Theorem 5.2, we know that the exponential bondage number of  $n$ -cycle is  $b_{exp}(C_n) = 1$  if  $n \equiv 0 \pmod{4}$  and  $b_{exp}(C_n) = 2$  otherwise. So, we get following corollaries.

**Corollary 5.3.** *If  $G$  is hamiltonian with  $n > 4$  vertices and  $\gamma_e(G) = \lceil \frac{n}{4} \rceil$ , then  $b_{exp}(G) \geq 2$  and in addition  $b_{exp}(G) \geq 1$  if  $n \equiv 0 \pmod{4}$ .*

**Corollary 5.4.** *If  $G$  is semihamiltonian with  $n > 4$  vertices and  $\gamma_e(G) = \lceil \frac{n}{4} \rceil$  and  $0 < n \equiv 0 \pmod{4}$ , then  $b_{exp}(G) \geq 2$ .*

**Theorem 5.5.** *The exponential bondage number of the star graph  $S_n$  of order ( $n \geq 3$ ) is  $b_{exp}(S_n) = 1$ .*

*Proof.* We know that  $\gamma_e(S_n) = 1$  by Theorem 2.2. The removal of an edge from the graph  $S_n$ , leaves a graph  $H$  consisting of an isolated vertex and star graph order  $n - 1$ . Thus,

$$\gamma_e(H) = 1 + \gamma_e(S_{n-1}) = 2 > \gamma_e(S_n).$$

Since  $\gamma_e(H) > \gamma_e(S_n)$  is obtained, the exponential bondage number of the star graph is  $b_{exp}(S_n) = 1$ . The proof is completed.  $\square$

**Theorem 5.6.** *The exponential bondage number of the complete graph of order  $n$  is  $b_{exp}(K_n) = \lceil \frac{n}{2} \rceil$ .*

*Proof.* Clearly,  $\gamma_e(K_n) = 1$  by Theorem 2.2. If any  $B_e$ -set  $F$  does not cover some vertex  $u$ , then  $\{u\}$  is a  $\gamma_e$ -set of  $K_n - F$ , so  $\gamma_e(K_n - F) = \gamma_e(K_n)$ , a contradiction. Hence  $F$  must cover every vertex, which implies  $|F| \geq \lceil \frac{n}{2} \rceil$ . Now define a specific  $F$  as follows. If  $n$  is even, let  $F$  be a matching of size  $\frac{n}{2}$ . If  $n$  is odd, let  $F$  consists of a matching of size  $\frac{n-1}{2}$  plus one vertex. So  $|F| = \lceil \frac{n}{2} \rceil$ . Let  $H = K_n - F$ . Then  $\gamma_e(H) \geq 2$  since  $F$  covers every vertex. So,  $F$  is a  $B_e$ -set. Consequently, the exponential bondage number of complete graph is  $b_{exp}(K_n) = \lceil \frac{n}{2} \rceil$ .

The proof is completed.  $\square$

As an immediate corollaries to Theorem 5.6, we have the following.

**Corollary 5.7.** *If  $K_n$  is a complete graph of order  $n$ , then  $b_{exp}(K_n) = \tau'(K_n)$ .*

**Corollary 5.8.** *If  $G$  is a connected graph of order  $n$ , then  $b_{exp}(G) \leq \lceil \frac{n}{2} \rceil$ .*

**Theorem 5.9.** *The exponential bondage number of the wheel graph of order  $n$  is  $b_{exp}(W_n) = 1$ .*

*Proof.* Let  $c$  be the central vertex of  $W_n$ . Since  $\deg(c) = n - 1$ , we have  $\gamma_e(W_n) = 1$  by Theorem 2.2. The removal of any edge which is incident to the vertex  $c$  in the graph  $W_n$  leaves a graph  $H$ . In the graph  $H$ , clearly  $\deg(c) = n - 2$ . Now we determine the exponential domination number of  $H$ . Let  $D$  be a  $\gamma_e$ -set of the graph  $H$ . If  $D = \{c\}$ , then  $D$  exponentially dominates  $n - 1$  vertices. Thus, there remains only one vertex does not exponentially dominated by  $D$ . Clearly, this vertex also must be in  $D$ . Then, we get  $\gamma_e(H) = 2$ .

Since  $\gamma_e(H) > \gamma_e(W_n)$ , the exponential bondage number of the wheel graph is  $b_{exp}(W_n) = 1$ . The proof is completed.  $\square$



## 6. CONCLUSION

In this study, a new graph theoretical parameter namely the exponential bondage number has been presented for the network vulnerability. The stability of popular interconnection networks has been studied and their exponential bondage numbers have been computed. These networks have been modeled with the complete graphs, the path graphs, the cycle graphs, the star graphs and the wheel graphs. Then upper bounds, lower bounds and exact formulas of the exponential bondage number have been obtained for any given graph  $G$ . As a further study, exact formulas or bounds may be obtained for graph operations and trees.

## REFERENCES

- [1] A.Aytac and B.Atay, *On Exponential Domination of Some Graphs*, Nonlinear Dynamics and Systems Theory **16(1)** (2016), 12-19.
- [2] A.Aytac and T.Turaci, *Vertex Vulnerability Parameter of Gear Graphs*, International Journal of Foundations of Computer Science **22(5)** (2011), 1187-1195.
- [3] A.Aytac, T.Turaci and Z.N.Odabas, *On The Bondage Number of Middle Graphs*, Mathematical Notes **93(6)** (2013), 803-811.
- [4] A.Aytac, Z.N.Odabas and T.Turaci, *The Bondage Number for Some Graphs*, Comptes Rendus de Lacademie Bulgare des Sciences **64(7)** (2011), 925-930.
- [5] C.A.Barefoot, B.Entringer and H.Swart, *Vulnerability in graphs-a comparative survey*, J. Combin. Math. Combin. Comput. **1** (1987), 13-22.
- [6] D.Bauer, F.Harary, J.Nieminen and C.L.Suffel, *Domination alteration sets in graph*, Discrete Math. **47** (1983), 153-161.
- [7] D.F.Hartnell and D.F.Rall, *Bounds on the bondage number of a graph*, Discrete Math. **128(1-3)** (1994), 173-177.
- [8] H.Frank and I.T.Frisch, *Analysis and design of survivable Networks*, IEEE Transactions on Communications Technology **18(5)** (1970), 501-519.
- [9] I.Mishkovski, M.Biey and L.Kocarev, *Vulnerability of complex Networks*, Commun. Nonlinear Sci Numer. Simulat. **16** (2011), 341-349.
- [10] J.F.Fink, M.S.Jacobson, L.F.Kinch and J.Roberts, *The bondage number of a graph*, Discrete Math. **86(1-3)** (1990), 47-57.
- [11] J.Huang and J.M.Xu, *The bondage number and efficient dominations of vertex-transitive graphs*, Discrete Math. **308** (2008), 571-582.
- [12] J.Kok and C.M.Mynhardt, *Reinforcement in Graphs*, Congr. Numer. **79** (1990), 225-231.
- [13] K.T.Newport and P.K.Varshney, *Design of survivable communication networks under performance constraints*, IEEE Transactions on Reliability **40** (1991), 433-440.
- [14] M.A.Henning, *Trees with Equal Average Domination and Independent Domination Numbers*, Ars Combin. **71** (2004), 305-318.
- [15] M.A.Henning and O.R.Oellermann, *The Average Connectivity of a Digraph*, Discrete App. Math. **140(1-3)** (2004), 143-153.
- [16] M.Blidia, M.Chellali and F.Maffray, *On Average Lower Independence and Domination Number in Graphs*, Discrete Math. **295** (2005), 1-11.
- [17] P.Dankelmann, D.Day, D.Erwin, S.Mukwembi and H.Swart, *Domination with Exponential Decay*, Discrete Math. **309** (2009), 5877-5883.
- [18] S.Skienna, *Implementing Discrete Mathematics: Combinatorics and Graph Theory with Mathematica*, Reading, MA: Addison-Wesley, 1990.
- [19] T.W.Haynes, S.T.Hedeniemi and P.J.Slater, *Fundamentals of Domination in Graphs*, Advanced Topic, Marcel Dekker, Inc, New York, 1998.
- [20] T.Turaci, *On the Average Lower Bondage Number of a Graph*, Rairo-Oper. Res. **50(4-5)** (2016), 1003-1012.

- [21] T.Turaci and E.Aslan, "The Average Lower Reinforcement Number of a Graph", *Rairo-Theor. Inf. Appl.* **50(2)** (2016), 135-144.
- [22] U.Teschner, *New results about the bondage number of a graph*, *Discrete Math.* **171** (1997), 249-259.
- [23] V.Aytac, *Average Lower Domination Number in Graphs*, *Comptes Rendus de Lacademie Bulgare des Sciences* **65(12)** (2012), 1665-1674.
- [24] V.Aytac and Z.N. Odabas, *On Computing the Vulnerability of Some Graphs As Average*, *International Journal of Pure and Applied Math.* **55(1)** (2009), 137-146.
- [25] V.Chvatal, *Tough graphs and Hamiltonian circuits*, *Discrete Math.* **5** (1973), 215-228.
- [26] W.K.Chiang and R.J.Chen, *The  $(n, k)$ -Star Graph: A Generalized Star Graph*, *Information Proc. Lett.* **56** (1995), 259-264.

DEPARTMENT OF COMPUTER ENGINEERING, FACULTY OF ENGINEERING, EGE UNIVERSITY 35100, IZMIR/TURKEY

*Email address:* `vecdi.aytac@ege.edu.tr`

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KARABUK UNIVERSITY 78050, KARABUK/TURKEY

*Email address:* `tufanturaci@karabuk.edu.tr`