

# AFFINOID SUBDOMAINS AS COMPLETIONS OF AFFINE SUBDOMAINS

GHIODEL GROZA

ABSTRACT. By following an idea of Nicolae Popescu, we construct affinoid subdomains as the completion of affine subdomains.

**Mathematics Subject Classification (2010):** 13B30, 18E35, 12J25, 54H13

**Keywords:** flat epimorphism of rings, affinoid algebra

*Article history:*

Received 16 June 2016

Received in revised form 28 June 2016

Accepted 30 June 2016

## 1. INTRODUCTION

Throughout this paper all rings are commutative with identity. Let  $A$  be a ring and let  $A[X_1, \dots, X_n]$  be the polynomial algebra over  $A$ . For simplicity, for any  $\nu = (i_1, \dots, i_n) \in \mathbb{N}^n$ , we denote  $\mathbf{X}^\nu = X_1^{i_1} \dots X_n^{i_n}$  and  $a_\nu = a_{i_1, \dots, i_n}$ . We also denote  $\mathbf{X} = (X_1, \dots, X_n)$  and  $N(\nu) = i_1 + i_2 + \dots + i_n$ . Thus we may write  $P \in A[\mathbf{X}]$  as

$$(1.1) \quad P = \sum_{\nu} a_{\nu} \mathbf{X}^{\nu}, \quad a_{\nu} \in A.$$

If  $g_1, \dots, g_n \in A$  and  $\nu = (i_1, \dots, i_n) \in \mathbb{N}^n$ , we denote  $\mathbf{g}^{\nu} = g_1^{i_1} \dots g_n^{i_n}$ .

Let  $A, B$  be two rings. A homomorphism of rings  $\phi : A \rightarrow B$  is called an *epimorphism of rings* if for any pair of homomorphisms of rings  $\psi_1, \psi_2 : B \rightarrow C$ , in another arbitrary ring  $C$ , the condition  $\psi_1 \phi = \psi_2 \phi$  implies  $\psi_1 = \psi_2$ . The epimorphism of rings  $\phi$  is called a *flat epimorphism of rings* if the  $A$ -module  $B$  is flat (see [1], Ch. 1).

The following result is known.

**Theorem 1.1.** ([4], p. 261) *Let  $\varphi : A \rightarrow B$  be a homomorphism of rings. The following assertions are equivalent:*

- a)  $\varphi$  is a flat epimorphism of rings.
- b) Let  $\mathcal{F} = \{I \text{ ideal of } A \text{ such that } \varphi(I)B = B\}$ . Then:
  - i) For any  $b \in B$ , there exists  $I \in \mathcal{F}$  such that  $\varphi(I)b \subseteq \varphi(A)$ ;
  - ii) If  $x \in A$ , and  $\varphi(x) = 0$ , there exists  $I \in \mathcal{F}$  such that  $Ix = 0$ .

If  $K$  is a field, a finitely generated  $K$ -algebra  $A$  is called an *affine  $K$ -algebra*. By an *affine subdomain* of  $\text{Sp } A := (\text{Max } A, A)$ , where  $\text{Max } A$  is the set of maximal ideals of  $A$ , we understand a subset  $\mathcal{U} \subset \text{Max } A$  and a homomorphism of affine algebras  $\varphi : A \rightarrow B$  such that:

- i)  $\varphi^a(\text{Max } B) \subset \mathcal{U}$ , where  $\varphi^a(M) := \varphi^{-1}(M)$ ,
- ii) If  $\psi : A \rightarrow C$  is a homomorphism of affine algebras such that  $\psi^a(\text{Max } C) \subset \mathcal{U}$ , then there exists a unique homomorphism of affine algebras  $\bar{\psi} : B \rightarrow C$  such that  $\bar{\psi} \phi = \psi$ .

Let  $A$  be a ring. A function  $\| \cdot \| : A \rightarrow [0, \infty)$  is called a *non-archimedean semi-norm* on  $A$  if the following properties are satisfied:

- i)  $\|0\| = 0$ ,
- ii)  $\|x - y\| \leq \max\{\|x\|, \|y\|\}$ , for all  $x, y \in A$ ,
- iii)  $\|xy\| \leq \|x\|\|y\|$ , for all  $x, y \in A$ ,
- iv)  $\|1\| \leq 1$ .

A non-archimedean semi-norm is called a *non-archimedean norm* if

- v)  $\|x\| = 0$ ,  $x \in A$ , implies  $x = 0$ .

In this case the pair  $(A, \|\cdot\|)$  is called a *normed ring*.

Let  $(A, \|\cdot\|_A)$  be a semi-normed ring (that is  $\|\cdot\|_A$  is a non-archimedean semi-norm on  $A$ ). If  $P \in A[X_1, \dots, X_n]$  is given by (1.1), define the *Gauss semi-norm* of  $P$  (see [2], p. 36) by

$$(1.2) \quad \|P\| = \max_{\nu} \|a_{\nu}\|_A.$$

Throughout this paper the semi-norm on  $A[X_1, \dots, X_n]$  will be the Gauss semi-norm.

If  $(A, \|\cdot\|)$  is a semi-normed ring and  $I$  be an ideal of  $A$ . Denote  $A/I$  the quotient ring of  $A$  with respect to  $I$  and  $\pi : A \rightarrow A/I$  the natural homomorphism. Then  $(A/I, \|\cdot\|_{\text{res}})$ , where

$$(1.3) \quad \|\pi(a)\|_{\text{res}} := \inf_{a' - a \in I} \|a'\|,$$

is a semi-normed ring. The corresponding topology on  $A/I$  is called the *quotient topology*.

Let  $A$  and  $B$  be two semi-normed rings. A ring homomorphism  $\phi : A \rightarrow B$  is said to be *strict* if the induced isomorphism  $\bar{\phi} : A/\text{Ker}\phi \rightarrow \phi(A)$  is a homeomorphism (see [2], p. 21). Here the topology on  $A/\text{Ker}\phi$  is the quotient topology and on  $\phi(A)$  we consider the induced topology from  $B$ .

If  $|\cdot|$  is a non-archimedean norm on  $A$  such that  $|xy| = |x||y|$ , for all  $x, y \in A$ , then  $|\cdot|$  is called a non-archimedean absolute value (valuation) on  $A$  and the pair  $(A, |\cdot|)$  is called a *valued ring*.

Let  $(K, |\cdot|)$  be a valued field and let  $A = K[X_1, \dots, X_n]/I$  be a  $K$ -affine algebra. Throughout this paper we consider on  $A$  the quotient topology defined by Gauss norm on  $K[X_1, \dots, X_n]$ .

Let  $(K, |\cdot|)$  be a complete valued field. For a positive integer  $n$  the following  $K$ -subalgebra of the  $K$ -algebra of formal power series in  $n$  indeterminates over  $K$  (see [2], p. 192):

$$T_n = K \langle X_1, \dots, X_n \rangle := \left\{ \sum_{i_1, \dots, i_n} a_{i_1 \dots i_n} X_1^{i_1} \dots X_n^{i_n} : a_{i_1 \dots i_n} \in K, \lim_{i_1 + \dots + i_n \rightarrow \infty} |a_{i_1 \dots i_n}| = 0 \right\}$$

is called the *Tate algebra in  $n$  indeterminates over  $K$* .

Each residue algebra  $T_n/I$  of  $T_n$  by an ideal  $I$  of  $T_n$  is a  $K$ -Banach algebra with respect to the residue norm defined by (1.3) (see [2], p. 221). This last  $K$ -Banach algebra  $T_n/I$  is called a  *$K$ -affinoid algebra*.

An *affinoid subdomain* of  $\text{Sp } A := (\text{Max } A, A)$ , where  $A$  is a  $K$ -affinoid algebra is a subset  $\mathcal{U} \subset \text{Max } A$  and a homomorphism of affinoid algebras  $\varphi : A \rightarrow B$  such that:

- i)  $\varphi^a(\text{Max } B) \subset \mathcal{U}$ , where  $\varphi^a(M) := \varphi^{-1}(M)$ ,
- ii) If  $\psi : A \rightarrow C$  is a homomorphism of affine algebras such that  $\psi^a(\text{Max } C) \subset \mathcal{U}$ , then there exists a unique homomorphism of affinoid algebras  $\bar{\psi} : B \rightarrow C$  such that  $\bar{\psi}\varphi = \psi$ .

As a corollary of a theorem of Gerritzen and Grauert (see [2], p. 309) it is known that an affinoid subdomain is a finite union of rational subdomains (defined in [2], p. 282). Moreover, a rational subdomain is constructed as the completion of a suitable ring of fractions (see [2], p. 232). As a continuation of the paper [3] my teacher Nicolae Popescu proposed, about ten years ago, to construct affinoid subdomains as completions of affine domains, which generalize the case when  $B$  is a ring of fractions of  $A$ . This paper, written to the *memory of Nicolae Popescu (1937-2010)*, is a first step in this direction.

The readers are expected to be familiar with the basic notations and results of commutative algebra and non-archimedean analysis, which can be found in, e.g. [5] and [2], respectively.

## 2. AFFINE SUBDOMAINS

Let  $A$  be a ring and let  $I = (g_1, g_2, \dots, g_n)$  be a finitely generated ideal of  $A$ . For a fixed non-negative integer  $m$ , denote

$$(2.1) \quad B = A[X_1, \dots, X_n]/J, \quad J = \left( \sum_{i=1}^n g_i X_i - 1, \mathbf{g}^\nu X_j - a_j^{(\nu)} \right),$$

where are considered all  $\nu = (i_1, \dots, i_n)$ , with  $N(\nu) = m$ , and  $a_j^{(\nu)} \in A$ ,  $j = 1, 2, \dots, n$ . Denote by  $\phi_I : A \rightarrow B$  the canonical homomorphism.

In order to give a sufficient condition under which  $\phi_I$  is a flat epimorphism of rings we prove the following result:

**Lemma 2.1.** *Let  $A$  be a ring and let  $m$  be a non-negative integer. If, in  $A[X_1, \dots, X_n]$ ,*

$$(2.2) \quad \sum_{\substack{\nu = (i_1, \dots, i_n) \\ N(\nu) \leq m}} a_\nu (\alpha_1 X_1 - \beta_1)^{i_1} \dots (\alpha_n X_n - \beta_n)^{i_n} = 0,$$

where  $a_\nu, \alpha_j, \beta_j \in A$ ,  $j = 1, 2, \dots, n$ , then for every  $\tau = (j_1, \dots, j_n)$ , with  $N(\tau) = m$  it follows that

$$(2.3) \quad \alpha^\tau a_\nu = 0, \text{ for all } \nu \text{ with } N(\nu) \leq m, \alpha = (\alpha_1, \dots, \alpha_n).$$

*Proof.* We use mathematical induction on  $m$ . Since (2.3) holds for  $m = 0$ , assume it holds for  $m = s$ .

We note that, for every  $\nu = (i_1, \dots, i_n)$ ,  $\delta = (j_1, \dots, j_n)$ , with  $N(\nu) = m$ ,  $N(\delta) \leq m - 1$ , there exist  $c_{\delta\nu} \in A$  such that

$$(2.4) \quad (\alpha_1 X_1 - \beta_1)^{i_1} \dots (\alpha_n X_n - \beta_n)^{i_n} = \alpha^\nu \mathbf{X}^\nu + \sum_{\substack{\delta = (j_1, \dots, j_n) \\ N(\delta) \leq m-1}} c_{\delta\nu} (\alpha_1 X_1 - \beta_1)^{j_1} \dots (\alpha_n X_n - \beta_n)^{j_n}.$$

Then, for  $m = s + 1$ , the equation (2.2) can be written as

$$(2.5) \quad \sum_{\substack{\tau = (j_1, \dots, j_n) \\ N(\tau) = s+1}} a_\tau \alpha^\tau \mathbf{X}^\tau + \sum_{\substack{\nu = (i_1, \dots, i_n) \\ N(\nu) \leq s}} a'_\nu (\alpha_1 X_1 - \beta_1)^{i_1} \dots (\alpha_n X_n - \beta_n)^{i_n} = 0,$$

where

$$(2.6) \quad a'_\nu = a_\nu + \sum_{\substack{\tau = (j_1, \dots, j_n) \\ N(\tau) = s+1}} a_\tau c_{\nu\tau}, \quad N(\nu) \leq s, \quad c_{\nu\tau} \in A.$$

By (2.5) we get

$$(2.7) \quad \alpha^\tau a_\tau = 0, \text{ for all } \tau = (j_1, \dots, j_n) \text{ with } N(\tau) = s + 1.$$

Since (2.3) holds for  $m = s$ , by equations (2.2), (2.5) and (2.7), it follows that for all  $\sigma = (r_1, \dots, r_n)$ , with  $N(\sigma) = s$ , we obtain

$$(2.8) \quad \alpha^\sigma a'_\nu = 0, \text{ for all } \nu \text{ with } N(\nu) \leq s.$$

Now, by (2.6)-(2.8), it follows that

$$(2.9) \quad \alpha^\tau a_\nu = 0, \text{ for all } \nu \text{ with } N(\nu) \leq s + 1,$$

which implies the lemma. □

**Theorem 2.2.** Let  $I = (g_1, \dots, g_n)$  be an ideal of  $A$  and let  $a_j^{(\nu)} \in A$ , where  $j = 1, 2, \dots, n$ ,  $N(\nu) = m$  and  $m$  is a fixed positive integer. If there exists  $N \in \mathbb{N}$  such that for all  $\tau$  with  $N(\tau) = m - 1$ ,

$$(2.10) \quad I^N(g^\tau - \sum_{j=1}^n a_j^{(\tau+\varepsilon^{(j)})}) = 0, \quad \varepsilon^{(j)} = (\delta_{1,j}, \dots, \delta_{n,j}),$$

$$(2.11) \quad I^N(a_j^{(\tau+\varepsilon^{(s)})} g_r - a_j^{(\tau+\varepsilon^{(r)})} g_s) = 0, \quad j, r, s = 1, 2, \dots, n,$$

then  $\phi_I : A \rightarrow B$ , where  $B$  is defined in (2.1), is a flat epimorphism of rings.

*Proof.* Let  $\mathcal{F} = \{I' : I' \text{ an ideal of } A, \varphi_{I'}(I)B = B\}$ . Then, by (2.1),  $I \in \mathcal{F}$  and, for all  $j = 1, 2, \dots, n$ ,  $\phi_I(I^m)\bar{X}_j \subset \phi_I(A)$ , where  $\bar{X}_j$  is the canonical image of  $X_j$  in  $B$ . Hence it follows that condition b) i) from Theorem 1.1 is fulfilled.

Now we verify condition b) ii) from Theorem 1.1.

If  $x \in A$  and  $\phi_I(x) = 0$ , then, for every  $j = 1, 2, \dots, n$  and  $\nu$ , with  $N(\nu) = m$ , there exist  $P, Q_j^{(\nu)} \in A[X_1, \dots, X_n]$  such that

$$(2.12) \quad x = P\left(\sum_{j=1}^n g_j X_j - 1\right) + \sum_{j=1}^n \sum_{\substack{\nu = (i_1, \dots, i_n) \\ N(\nu) = m}} Q_j^{(\nu)}(\mathbf{g}^\nu X_j - a_j^{(\nu)}).$$

If  $\sigma = (r_1, \dots, r_n)$ , with  $N(\sigma) = m$ , there exists a positive integer  $t$ , and  $\tau$  with  $N(\tau) = m - 1$  such that  $\sigma = \tau + \varepsilon^{(t)}$ . Hence, by (2.10),  $I^N g^\sigma = I^N g^{\tau+\varepsilon^{(t)}} = I^N \sum_{j=1}^n a_j^{(\tau+\varepsilon^{(j)})} g_t$  and

$$(2.13) \quad I^N \left( \mathbf{g}^\sigma x - P \sum_{j=1}^n (\mathbf{g}^\sigma g_j X_j - g_t a_j^{(\tau+\varepsilon^{(j)})}) - \mathbf{g}^\sigma \sum_{j=1}^n \sum_{\substack{\nu = (i_1, \dots, i_n) \\ N(\nu) = m}} (\mathbf{g}^\nu X_j - a_j^{(\nu)}) Q_j^{(\nu)} \right) = 0.$$

Since, by (2.11),  $I^N(g_t a_j^{(\tau+\varepsilon^{(j)})} - g_j a_j^{(\sigma)}) = 0$  and  $I^N(\mathbf{g}^\sigma a_j^{(\nu)} - \mathbf{g}^\nu a_j^{(\sigma)}) = 0$ , by denoting

$$(2.14) \quad S_j = g_j P + \sum_{\substack{\nu = (i_1, \dots, i_n) \\ N(\nu) = m}} \mathbf{g}^\nu Q_j^{(\nu)},$$

the equation (2.13) becomes

$$(2.15) \quad I^N \left( \mathbf{g}^\sigma x - \sum_{j=1}^n (\mathbf{g}^\sigma X_j - a_j^{(\sigma)}) S_j \right) = 0, \quad \text{for all } \sigma \text{ with } N(\sigma) = m.$$

Denote

$$d = \max_{1 \leq j \leq n} (\deg S_j),$$

which, by (2.14), is independent of  $\nu$ . Then, by (2.4), for all  $\theta = (s_1, \dots, s_n)$ , with  $N(\theta) = ndm + 1$ , it follows that there exists  $\sigma = (r_1, \dots, r_n)$ , with  $N(\sigma) = m$ , such that

$$(2.16) \quad \mathbf{g}^\theta S_j = \sum_{\substack{\delta = (t_1, \dots, t_n) \\ N(\delta) \leq d}} b_j^{(\delta)} \left( \mathbf{g}^\sigma X_1 - a_1^{(\sigma)} \right)^{t_1} \dots \left( \mathbf{g}^\sigma X_n - a_n^{(\sigma)} \right)^{t_n}, \quad b_j^{(\delta)} \in A.$$

Since  $I^N$  is finitely generated, by (2.15), (2.16) and Lemma 2.1 with  $a_0 = \mathbf{g}^{\theta+\gamma}$ , where  $N(\gamma) = N$ , it follows that  $I^M x = 0$ , where  $M \geq N + m + ndm + 1$ . Thus the condition b) ii) from Theorem 1.1 holds and  $\phi_I$  is a flat epimorphism of rings.  $\square$

**Corollary 2.3.** *Under the hypotheses of Theorem 2.2, for all  $I_1 \in \mathcal{F}$ , there exists a non-negative integer  $M$  such that  $I^M \subset I_1$ .*

*Proof.* If  $I_1 \in \mathcal{F}$ , there exist a positive integer  $t$ ,  $x_i \in I_1$ ,  $b_i \in B$ ,  $i = 1, 2, \dots, t$ , such that

$$(2.17) \quad \sum_{i=1}^t \varphi_I(x_i) b_i = 1.$$

By Theorems 1.1 b) i) and 2.2 we can choose a non-negative integer  $M_1$  such that  $\varphi_I(I^{M_1}) b_i \subset \varphi_I(A)$ . Hence we get, for all  $\sigma$ , with  $N(\sigma) = M_1$ ,

$$(2.18) \quad \varphi_I(\mathbf{g}^\sigma) b_i = \varphi_I(\alpha_i^{(\sigma)}), \quad \alpha_i^{(\sigma)} \in A.$$

By (2.17) and (2.18) it follows that

$$\varphi_I(\mathbf{g}^\sigma) = \sum_{i=1}^t \varphi_I(x_i \alpha_i^{(\sigma)}),$$

and by Theorem 2.2 and by Theorem 1.1 b) ii) there exists a non-negative integer  $M_2$  such that, for all  $\sigma$ , with  $N(\sigma) = M_1$ , we get

$$(2.19) \quad I^{M_2}(\mathbf{g}^\sigma - \sum_{i=1}^t x_i \alpha_i^{(\sigma)}) = 0.$$

Since  $x_i \in I_1$ , by (2.19), it follows that for  $M = M_1 + M_2$ ,  $I^M \subset I_1$ .  $\square$

**Example 2.4.** Let  $A$  be a ring and let  $I = (g_1, \dots, g_n)$  be an ideal of  $A$ . We choose, for example, the elements  $b_j^{(s)} \in A$ ,  $j, s = 1, 2, \dots, n$ , such that  $\sum_{j=1}^n b_j^{(j)} = 1$ , and, for  $j \neq s$ ,  $b_j^{(s)} = g_s$ . If we take  $a_j^{(\tau+\varepsilon^{(s)})} = \mathbf{g}^\tau b_j^{(s)}$ , it follows that (2.10) and (2.11) hold. Thus  $\varphi_I$  is a flat epimorphism of rings.

**Remark 2.5.** Let  $K$  be a field and let  $A$  be a  $K$ -affine algebra. If  $B$  is defined by (2.1), then, by Theorem 2.2,  $\mathcal{U} = \phi_I^q(\text{Max } B)$ , is an affine subdomain of  $\text{Sp } A = (\mathcal{U}, A)$  (see [3]).

**Theorem 2.6.** *Let  $K$  be a field and let  $\phi : A \rightarrow B$  be a homomorphism of  $K$ -affine algebras such that  $\mathcal{U} = \phi^q(\text{Max } B)$  and  $\phi$  define an affine subdomain of  $\text{Sp } A$ . Let  $\mathcal{F} = \{I' \text{ ideal in } A; \phi(I')B = B\}$ . If there exists  $I \in \mathcal{F}$  such that, for all  $I' \in \mathcal{F}$ , there exists a positive integer  $t$  such that  $I^t \subset I'$ , then there exist the positive integers  $n, N, m$ , and for all  $\tau \in \mathbb{N}^n$  with  $N(\tau) = m - 1$ ,  $i = 1, 2, \dots, n$ , there exist  $a_i^{(\tau+\varepsilon^{(s)})} \in A$ ,  $s = 1, 2, \dots, n$ , such that we can take  $I = (g_1, \dots, g_n)$  such that (2.10), (2.11) hold.*

*Proof.* Since  $\mathcal{U}$  and  $\phi$  define an affine subdomain of  $\text{Sp } A$ , by Theorem 3.2 from [3],  $\phi$  is a flat epimorphism of rings. Because  $I \in \mathcal{F}$  it follows that there exists a positive integer  $n$  such that

$$(2.20) \quad \sum_{i=1}^n \phi(g_i) b_i = 1, \quad g_i \in I, \quad b_i \in B.$$

Without loss of generality we may assume  $I = (g_1, \dots, g_n)$ . By Theorem 1.1 b) i) and by hypotheses there exists a positive integer  $m$  such that, for all  $\nu$  with  $N(\nu) = m$ , we get

$$(2.21) \quad \phi(g^\nu) b_i = \phi(a_i^{(\nu)}), \quad a_i^{(\nu)} \in A.$$

If  $\tau \in \mathbb{N}^n$  with  $N(\tau) = m - 1$ , by (2.21),

$$\phi(a_i^{(\tau+\varepsilon^{(r)})}) \phi(g_s) = \phi(g^{\tau+\varepsilon^{(r)}}) b_i \phi(g_s) = \phi(g^{\tau+\varepsilon^{(s)}}) b_i \phi(g_r) = \phi(a_i^{(\tau+\varepsilon^{(s)})}) \phi(g_r), \quad r, s = 1, \dots, n.$$

Then, by Theorem 1.1 b) ii), there exists a positive integer  $n_1$  such that

$$I^{n_1}(a_j^{(\tau+\varepsilon^{(s)})}g_r - a_j^{(\tau+\varepsilon^{(r)})}g_s) = 0, \quad j, r, s = 1, 2, \dots, n.$$

Similarly, by (2.20) and (2.21), we get

$$\phi(g^\tau) = \sum_{j=1}^n \phi(g^{\tau+\varepsilon^{(j)}})b_j = \sum_{j=1}^n \phi(a_j^{(\tau+\varepsilon^{(j)})}).$$

Then, by Theorem 1.1 b) ii), there exists a positive integer  $n_2$  such that

$$I^{n_2}(g^\tau - \sum_{j=1}^n a_j^{(\tau+\varepsilon^{(j)})}) = 0.$$

By taking  $N = \max\{n_1, n_2\}$  it follows the statement of the theorem.  $\square$

### 3. AFFINOID SUBDOMAINS

Let  $K$  be a complete non-archimedean valued field and let  $A$  be a  $K$ -affine algebra. We need the following result:

**Lemma 3.1.** *Let  $A = K[Z_1, \dots, Z_r]/I_1$  be a  $K$ -affine algebra, where  $I_1$  is an ideal of  $K[Z_1, \dots, Z_r]$ . Then  $\tilde{A}$  (the completion of  $A$  with respect to the residue semi-norm defined by Gauss semi-norm) is an affinoid  $K$ -algebra.*

*Proof.* Since the canonical homomorphism of semi-normed  $K$ -affine algebra  $\pi_A : K[Z_1, \dots, Z_r] \rightarrow A$  is a strict homomorphism which is onto, by Corollary 6 from [2], p. 23, we get that  $\tilde{\pi}_A : K \langle Z_1, \dots, Z_r \rangle \rightarrow \tilde{A}$  is onto. Hence it follows the lemma.  $\square$

If  $I$  is an ideal of  $A$ , denote by  $A_I$  the algebra  $B$  defined in (2.1).

**Theorem 3.2.** *Let  $K$  be a complete non-archimedean valued field, let  $A$  be a  $K$ -affine algebra and let  $I$  be an ideal of  $A$  satisfying the conditions (2.10) and (2.11) (see Theorem 2.6). Then the canonical homomorphism  $\tilde{\phi}_I : \tilde{A} \rightarrow \tilde{A}_I$  defines the affinoid subdomain  $\mathcal{U} = \tilde{\phi}_I^a(\text{Max } \tilde{A}_I)$  of  $\text{Sp } \tilde{A}$ .*

*Proof.* By the canonical commutative diagram

$$\begin{array}{ccc} A[X_1, \dots, X_n] & \xrightarrow{\pi} & A_I \\ \downarrow i_{A[X_1, \dots, X_n]} & & \downarrow i_{A_I} \\ \tilde{A} \langle X_1, \dots, X_n \rangle & \xrightarrow{\tilde{\pi}} & \tilde{A}_I \end{array} ,$$

where  $\pi$  is a strict homomorphism of rings which is onto and, by Proposition 5 from [2], p. 22, it follows that  $\tilde{A}_I \cong \tilde{A} \langle X_1, \dots, X_n \rangle / J\tilde{A} \langle X_1, \dots, X_n \rangle$ , because  $\tilde{J} = J\tilde{A} \langle X_1, \dots, X_n \rangle$  (see [2], Proposition 3, p. 222).

Let  $\psi : \tilde{A} \rightarrow C$  be a homomorphism of  $K$ -affinoid algebras such  $\psi^a(\text{Max } C) \subset \tilde{\phi}_I^a(\text{Max } \tilde{A}_I)$ . We prove that  $\psi(I)C = C$ .

Suppose the contrary. Then there exists  $M_C \in \text{Max } C$  such that  $\psi(I)C \subset M_C$ . Hence  $I \subset \psi^a(M_C) = \tilde{\phi}_I^a(M)$ , where  $M \in \text{Max } \tilde{A}_I$ . Then  $\tilde{\phi}_I(I) \subset M$ , a contradiction since  $\phi_I(I)A_I = A_I$  implies  $\tilde{\phi}_I(I)\tilde{A}_I = \tilde{A}_I$ . Thus  $\psi(I)C = C$  and there exist  $d^{(1)}, \dots, d^{(n)} \in C$  such that

$$(3.1) \quad \sum_{i=1}^n \psi(g_i)d^{(i)} = 1.$$

We identify  $\tilde{A}_I$  with  $\tilde{A} \langle X_1, \dots, X_n \rangle / J\tilde{A} \langle X_1, \dots, X_n \rangle$ , and, by considering  $c^{(i)} = \bar{X}_i$ , from (2.1) we get

$$(3.2) \quad \sum_{i=1}^n \tilde{\phi}_I(g_i) c^{(i)} = 1$$

and

$$(3.3) \quad \tilde{\phi}_I(\mathbf{g}^\nu) c^{(i)} = \tilde{\phi}_I(a_i^\nu), \quad i = 1, 2, \dots, n, \quad \text{for all } \nu \text{ with } N(\nu) = m.$$

For an arbitrary positive integer  $r$ , by (3.1), it follows that

$$(3.4) \quad \sum_{\sigma; N(\sigma)=r}^n \psi(\mathbf{g}^\sigma) d^{(\sigma)} = 1,$$

where  $d^{(\sigma)}$  are monomials of degree  $r$  in  $d^{(1)}, \dots, d^{(n)}$  whose coefficients are non-negative integers.

By multiplying (3.1) by  $\psi(a_j^{(\tau+\varepsilon^{(j)})} \mathbf{g}^\delta)$ , where  $N(\tau) = m - 1$ ,  $N(\delta) = N$  and by using (2.11) we find

$$\psi(\mathbf{g}^\delta) \psi(a_j^{(\tau+\varepsilon^{(j)})}) = \sum_{i=1}^n \psi(a_j^{(\tau+\varepsilon^{(i)})}) \psi(g_j) d^{(i)} \psi(\mathbf{g}^\delta).$$

By multiplying by  $d^{(\delta)}$ , by summing with respect to  $\delta$ , with  $N(\delta) = N$ , and by using (3.4) we get

$$(3.5) \quad \psi(a_j^{(\tau+\varepsilon^{(j)})}) = \sum_{i=1}^n \psi(a_j^{(\tau+\varepsilon^{(i)})}) d^{(i)} \psi(g_j), \quad \text{for all } \tau \text{ with } N(\tau) = m - 1.$$

By multiplying (3.5) by  $\psi(\mathbf{g}^\delta)$ , by summing with respect to  $j$ , and by using (2.10) it follows that

$$\psi(\mathbf{g}^\delta) \psi(\mathbf{g}^\tau) = \psi(\mathbf{g}^\delta) \sum_{j=1}^n \sum_{i=1}^n \psi(a_j^{(\tau+\varepsilon^{(i)})}) d^{(i)} \psi(g_j).$$

Then, by multiplying once again by  $d^{(\delta)}$  and by summing with respect to  $\delta$ , we find

$$(3.6) \quad \psi(\mathbf{g}^\tau) = \sum_{j=1}^n \sum_{i=1}^n \psi(a_j^{(\tau+\varepsilon^{(i)})}) d^{(i)} \psi(g_j).$$

By multiplying (3.6) by  $d^{(\tau)}$ , with  $N(\tau) = m - 1$ , and, by using (3.4), we get

$$(3.7) \quad \sum_{j=1}^n \left( \sum_{\tau, N(\tau)=m-1} \sum_{i=1}^n \psi(a_j^{(\tau+\varepsilon^{(i)})}) d^{(i)} d^{(\tau)} \right) \psi(g_j) = 1.$$

If we denote, for  $j = 1, 2, \dots, n$ ,

$$(3.8) \quad \tilde{d}^{(j)} = \sum_{\tau, N(\tau)=m-1} \sum_{i=1}^n \psi(a_j^{(\tau+\varepsilon^{(i)})}) d^{(i)} d^{(\tau)},$$

then, from (3.7), we find

$$(3.9) \quad \sum_{j=1}^n \psi(g_j) \tilde{d}^{(j)} = 1.$$

If  $N(\nu) = m$ ,  $N(\delta) = N$ , by (2.11) and (3.8), it follows that

$$\psi(\mathbf{g}^{\nu+\delta}) \tilde{d}^{(j)} = \sum_{\tau, N(\tau)=m-1} \sum_{i=1}^n \psi(a_j^{(\tau+\varepsilon^{(i)})}) d^{(i)} d^{(\tau)} \psi(\mathbf{g}^\nu) \psi(\mathbf{g}^\delta)$$

$$\begin{aligned}
&= \psi(\mathbf{g}^\delta) \sum_{\tau, N(\tau)=m-1} \sum_{i=1}^n \psi(a_j^{(\nu)}) \psi(\mathbf{g}^{\tau+\varepsilon^{(i)}}) d^{(\tau)} d^{(i)} \\
&= \psi(\mathbf{g}^\delta) \psi(a_j^{(\nu)}) \sum_{\tau, N(\tau)=m-1} \sum_{i=1}^n \psi(\mathbf{g}^\tau) d^{(\tau)} \psi(\mathbf{g}^{\varepsilon^{(i)}}) d^{(i)} = \psi(\mathbf{g}^\delta) \psi(a_j^{(\nu)}).
\end{aligned}$$

Hence

$$\psi(\mathbf{g}^\delta) \psi(\mathbf{g}^\nu) \tilde{d}^{(j)} = \psi(\mathbf{g}^\delta) \psi(a_j^{(\nu)}).$$

By multiplying by  $d^{(\delta)}$  and by summing with respect to  $\delta$ , we find

$$(3.10) \quad \psi(\mathbf{g}^\nu) \tilde{d}^{(j)} = \psi(a_j^{(\nu)}), \quad j = 1, 2, \dots, n.$$

Let  $M_C \in \text{Max } C$ . Then  $C/M_C$  is a finite extension of  $K$  (see [2], Corollary 3, p. 228) and

$$(3.11) \quad |\psi(\mathbf{g}^\nu)|_{C/M_C} = |\mathbf{g}^\nu|_{\tilde{A}/\psi^a(M_C)} = |\mathbf{g}^\nu|_{\tilde{A}/\tilde{\phi}_I^a(M)} = |\tilde{\phi}_I(\mathbf{g}^\nu)|_{\tilde{A}_I/M},$$

where  $M \in \text{Max } \tilde{A}_I$ ,  $|\cdot|_{C/M_C}$  is the unique absolute value on  $C/M_C$  which extends the absolute value on  $K$  and  $\psi^a(M_C) = \tilde{\phi}_I^a(M)$  (see [2]).

Similarly we get

$$(3.12) \quad |\psi(a_j^{(\nu)})|_{C/M_C} = |\tilde{\phi}_I(a_j^{(\nu)})|_{\tilde{A}_I/M}.$$

By (3.3), (3.10)-(3.12) it follows that, for all  $M_C \in \text{Max } C$ ,

$$(3.13) \quad |\tilde{d}^{(j)}|_{C/M_C} = |c^{(j)}|_{\tilde{A}_I/M}.$$

Hence (see [2], p. 169 and p. 236)

$$(3.14) \quad \|\tilde{d}^{(j)}\|_{\text{sup}} \leq |c^{(j)}|_{\text{sup}} \leq 1,$$

and the elements  $\tilde{d}^{(j)}$  are power bounded (see [2], Proposition 1, p. 240). Then, by using Proposition 4 from [2], p. 222, there exists a continuous mapping  $\theta_{\tilde{A}} : \tilde{A} < X_1, \dots, X_n > \rightarrow C$  such that

$$\theta_{\tilde{A}}(X_j) = \tilde{d}^{(j)} \text{ and } \theta_{\tilde{A}}/\tilde{A} = \psi.$$

By (3.9) and (3.10) we get  $J\tilde{A} < X_1, \dots, X_n > \subset \text{Ker } \theta_{\tilde{A}}$ . Thus there exists a continuous mapping  $\theta : \tilde{A}_I \rightarrow C$  such that

$$(3.15) \quad \theta_{\tilde{A}_I} = \psi.$$

If  $\theta'_{\tilde{A}_I} = \theta_{\tilde{A}_I}$ , because  $\tilde{\phi}_I i_A = i_{A_I} \phi_I$ , and  $\phi_I$  is an epimorphism of rings, it follows that  $\theta'_{i_{A_I}} = \theta_{i_{A_I}}$ . Since  $i_{A_I}(A_I)$  is dense in  $\tilde{A}_I$  we get  $\theta' = \theta$ . Hence  $\tilde{A}_I$  is an affinoid subdomain of  $\text{Sp } \tilde{A}$ .  $\square$

#### REFERENCES

- [1] N. Bourbaki, *Éléments de mathématique. Algèbre commutative*, Ch. I-II, 1961, Hermann Paris.
- [2] S. Bosch, U. Günter, R. Remmert, *Non-Archimedean Analysis*, Springer-Verlag, Berlin, 1984.
- [3] G. Groza, N. Popescu, On affine subdomains, *Rev. Roum. Math. Pures Appl.*, **49**(2004), 3, 231-246.
- [4] N. Popescu, *Abelian Categories with Applications to Rings and Modules*, L.M.S. Monographs, Academic Press, London and New-York, 1973.
- [5] O. Zariski, P. Samuel, *Commutative Algebra*, Vol. 1, Springer-Verlag, New York, 1958.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, TECHNICAL UNIVERSITY OF CIVIL ENGINEERING BUCHAREST, 020396, ROMANIA,

*E-mail address:* grozag@utcb.ro